



GC-MS ANALYSIS OF MICROORGANISM MARKERS IN PLANTS (REVIEW)

Bahranova M.A., Mukhamadiev
N.K., Khalilov K.F., Mukhamadiev
A.N., Alikulov B.S.

Received 15th Aug 2023,
Accepted 26th Sep 2023,
Online 5th Oct 2023

Samarkand State University named
after Sharof Rashidov, Samarkand,
Republic of Uzbekistan E- mail :
baxranovamohidil@gmail.com

ABSTRACT *The presence of microorganisms in plants can have a significant impact on their growth, development and general health. Traditional methods for identifying and characterizing these microorganisms can be time consuming and often lack specificity. In recent years, gas chromatography-mass spectrometry (GC-MS) has become a powerful method for the analysis of microorganism markers in plants. This review article presents the principles and applications of GC-MS in the analysis of microorganism markers, indicating its advantages and limitations. In addition, it discusses various case studies in which GC-MS has been successfully used to detect and identify microorganisms in plants, paving the way for improved plant disease control strategies.*

Keywords *microorganism, marker, fatty acids, gas chromatography, analysis.*

Relevance:

Plants are known to contain a wide range of microorganisms, including bacteria, fungi, and viruses, which can have both positive and negative effects on plant health [1-3]. The identification and characterization of these microorganisms is critical to understanding their role in plant-microbial interactions. GC-MS, with its high sensitivity and the ability to obtain detailed chemical information, has become a valuable tool in this field. In this regard, for completeness of information, it is necessary to consider the following sections:

Principles of GC-MS. GC/MS principles include the following [4-8]:

1. **Separation of a mixture of components:** GC-MS is based on the separation of a mixture of analytes into individual components using gas chromatography. This is achieved due to the different affinities of the components for the stationary phase (stationary column) and the mobile phase (mobile gas).

2. **Ionization of the components:** After the separation of the components, they are ionized, which allows them to turn into ions. Various methods are used for this, such as electron ionization (EI), chemical ionization (CI) or electronic shielding (EI).

3. **Mass analysis ionized components:** The ionized components pass through a mass analyzer, which separates the ionized particles by mass and charge. Typically, GC/MS uses mass filters such as a quadrupole or time line device (TOF).

4. **Detection and registration of ions:** Ions are registered by a detector and further processed using a mass spectrometer. This allows you to determine the mass and relative content of each component in the sample.

5. **Component Identification:** To identify components, mass spectrum databases are used, which contain information on the mass spectra of various substances. By comparing the resulting mass spectrum with databases, the structure can be determined and analytes identified.

These principles form the basis of GC-MS analysis and allow high accuracy and sensitivity in the determination of components in various samples.

Microorganism markers in plants. This section discusses the different types of microorganism markers that can be analyzed using GC/MS. It covers volatile organic compounds (VOCs), non-volatile metabolites, fatty acids, and other biomarkers that can serve as indicators of the presence of microbes in plants.

Sample preparation methods in GC-MS.

Sample preparation methods in gas chromatography -mass spectrometry (GC-MS) may vary depending on the type of sample being analyzed and the information required.

The following are some of the main methods for sample preparation in GC/MS:

➤ **Extraction:** This method is used to extract analytes from the sample matrix. For this, various solvents are used, such as acetonitrile, chloroform, diethyl ether, etc. The extraction process can be carried out using ultrasonic treatment or cold irrigation.

➤ **Evaporation:** In this method, the sample is subjected to heat treatment to concentrate the analytes. Heating can be carried out in a water bath, steam bath, vacuum film evaporator or other special devices.

➤ **Direct Injection:** This method involves directly injecting the sample into the GC/MS for analysis. The sample, usually dissolved in a solvent, is injected through the apparatus into the gas chromatograph column.

➤ **Derivatives:** Sometimes analytes can be chemically modified to improve their stability or detectability. Some common derivative methods include derivatization amine or carbonyl groups, the conversion of alkanes into gasoline or aromatic compounds, and others.

➤ **Fractionation:** This method is used when the sample contains many analytes with different physicochemical properties. In such cases, the sample may be fractionated based on various filtration, extraction or chromatography techniques.

The choice of the appropriate GC/MS sample preparation method depends on the purpose of the analysis, the type of sample and its matrix, and the capabilities and limitations of the GC/MS setup. It is important to take into account the specific requirements of each assay and tailor sample preparation methods to meet these requirements.

Application of GC-MS in the analysis of microorganisms. This section provides a comprehensive overview of the use of GC-MS in the analysis of microorganism markers in plants. It discusses case studies where GC-MS has been successfully used to detect and identify specific microorganisms such as fungal pathogens, bacterial endophytes, and viral infections. In addition, it highlights the use of GC-MS to study plant-microbial interactions and the role of micro-organisms in plant diseases.

The importance of GC-MS in the analysis of microbial markers in plants cannot be overestimated. Its ability to provide detailed chemical information about the metabolites of microorganisms provides valuable insight into the complex relationships between plants and microorganisms. Using the power of GC/MS, researchers can unravel the mysteries of plant-microbial interactions, leading to the development of sustainable and effective strategies to protect plant health in the face of microbial challenges.

Microorganisms can leave various markers or signatures that can be used for their detection and identification [9-12]. Here are some commonly studied microorganism markers (Table 1):

Microorganism markers	➤ Volatile organic compounds
	➤ Fatty acid
	➤ Metabolites
	➤ Cell wall components
	➤ Nucleic acids
	➤ Proteins and enzymes
	➤ Biomolecules and metabolic pathways

1. Volatile Organic Compounds (VOC): Microorganisms can produce a wide range of volatile compounds that can serve as markers. These include, among others, alcohols, aldehydes, ketones, esters and sulfur compounds. VOCs are often responsible for characteristic odors associated with microbial growth [13–16].

2. Fatty acids: The fatty acid composition of microorganisms may vary by species and strain. Fatty acid profiles can be used as markers for the identification and classification of microorganisms [17-21].

3. Metabolites: Microorganisms produce specific metabolites during their metabolic activity. These metabolites, such as secondary metabolites, can serve as markers of microbial presence and activity [22–25]. Examples include antibiotics, mycotoxins and pigments.

4. Cell wall components: The composition and structure of cell wall components can vary between microorganisms, providing characteristic markers for their identification [26-29]. For example, peptidoglycan is a major component of bacterial cell walls.

5. Nucleic acids: The genetic material of microorganisms, such as DNA and RNA, can serve as a marker for their detection and identification. Certain gene sequences or regions can be targeted using molecular techniques such as polymerase chain reaction (PCR) for microbial identification [30–34].

6. Proteins and enzymes. Microorganisms express unique proteins and enzymes that can act as markers. These markers can be targeted using immunological methods such as enzyme-linked immunosorbent assay (ELISA) to detect and identify microorganisms [35-38].

7. Biomolecules and metabolic pathways. Microorganisms have different metabolic pathways and produce specific biomolecules. They may include amino acids, sugars, organic acids, and enzymes that can be used as markers for the presence and activity of microbes.

It is important to note that microorganism markers may vary depending on the specific species, strain, and environmental conditions. Researchers often use a combination of markers and analytical methods such as gas chromatography-mass spectrometry (GC-MS) to achieve accurate and reliable identification and characterization of microorganisms in a variety of samples.

Fatty acids are commonly used as markers for the identification and classification of microorganisms. The composition and relative amount of fatty acids can vary among different species and strains of microbes, making them useful for distinguishing microorganisms. Here are some examples of fatty acids that act as microbial markers (Table 2):

Fatty acid markers	➤ Branched-chain fatty acids (BCFAs)
	➤ Monounsaturated fatty acids (MUFAs)

➤ Polyunsaturated fatty acids (PUFAs)
➤ Cyclopropane fatty acids (CFA)
➤ Hydroxy fatty acids
➤ Saturated fatty acids (SFAs)
➤ Odd chain fatty acids
➤ Short chain fatty acids (SCFA)
➤ Long chain fatty acids (LCAs)
➤ Omega-3 fatty acids
➤ Hydroxylated fatty acids
➤ Polyhydroxyalkanoates (PHA)
➤ Omega 9 fatty acids
➤ conjugated fatty acids
➤ Iso/ anteiso fatty acids
➤ 3-hydroxy fatty acids
➤ Mycolic acids
➤ Gopanoids
➤ Wax esters
➤ Sterols
➤ Fatty acid phospholipids (PLFA)
➤ Lipopolysaccharides (LPS)
➤ Essential lipids
➤ Very long chain fatty acids (VLCFA)
➤ diacylglycerols

1. Branched Chain Fatty Acids (BCFAs): BCFAs are commonly found in bacterial cell membranes and can be used as markers to identify bacteria. Some examples of BCFAs include o- and anteiso -fatty acids such as iso-C15:0 and iso-C17:0 [13,15,39-44].

2. Monounsaturated fatty acids (MUFAs): MUFAs are fatty acids with one double bond in their hydrocarbon chain. Microorganisms often display specific MUFA patterns that can be used for microbial identification. Examples include oleic acid (C18:1) and palmitoleic acid (C16:1) [13,39,45-47].

3. Polyunsaturated fatty acids (PUFAs): PUFAs are fatty acids with several double bonds in their hydrocarbon chain. Although the presence of certain PUFAs in microorganisms is less common than in higher organisms, they may be indicative of certain types of microbes. Examples include linoleic acid (C18:2) and linolenic acid (C18:3) [48-51].

4.Cyclopropane fatty acids (CFAs): Some bacteria produce CFAs, which are fatty acids with one or more cyclopropane rings in their structure. The presence of CFAs can be a characteristic marker for certain bacterial species. Cyclopropane fatty acids such as cyclopropane-C19:0 are commonly used for microbial identification [15,39,49,52-54].

5.Hydroxy fatty acids: Hydroxy fatty acids are fatty acids that contain one or more hydroxyl groups in their structure. These fatty acids are often produced by certain types of bacteria and can serve as markers for their identification [55-58]. Examples include 3-hydroxy fatty acids such as 3-hydroxydecanoic acid (C10:0 3-OH) and 3-hydroxytetradecanoic acid (C14:0 3-OH).

6. Saturated fatty acids (SFA). Saturated fatty acids are fatty acids that do not have double bonds in the hydrocarbon chain. Although SFAs are not as specific as other types of fatty acids, their relative abundance and distribution can provide insight into microbial communities [14,47,49]. Examples of SFAs include stearic acid (C18:0) and palmitic acid (C16:0).

7. Odd-chain fatty acids. Odd chain fatty acids have an odd number of carbon atoms in their structure, such as heptadecanoic acid (C17:0) and nonadecanoic acid (C19:0). These fatty acids can be produced by some microorganisms and can serve as markers for their identification [15,60-64].

8.Short-Chain Fatty Acids (SCFAs): SCFAs typically contain less than six carbon atoms and are produced by microbial fermentation. Examples include acetic acid (C2:0) and propionic acid (C3:0). SCFAs play an important role in microbial ecology and can be used as markers of specific microbial metabolic activity [65–68].

9. Long Chain Fatty Acids (LCFAs): LCFAs are fatty acids with more than 14 carbon atoms. They can be found in a variety of microorganisms, and their relative abundance can provide insight into the composition of a microbial community. Examples include behenic acid (C22:0) and arachidic acid (C20:0) [13,14,64].

10. Omega-3 fatty acids: Omega-3 fatty acids are polyunsaturated fatty acids with a double bond at the third carbon atom from the omega end of the carbon chain. They are commonly found in some microorganisms, especially marine bacteria and algae. Examples of omega-3 fatty acids include eicosapentaenoic acid (EPA, C20:5 ω 3) and docosahexaenoic acid (DHA, C22:6 ω 3). The presence of omega-3 fatty acids can serve as a marker for certain types or groups of microbes [69-74].

11. Hydroxylated fatty acids: Hydroxylated fatty acids are fatty acids that contain one or more hydroxyl groups attached to the carbon chain. These fatty acids are often produced by some bacteria as secondary metabolites and may act as signaling molecules or play a role in host-microbe interactions. Examples include 10-hydroxy-2-decanoic acid (C10:0 10-OH) produced by some Gram-negative bacteria [13,75-78].

12.Polyhydroxyalkanoates (PHA): Polyhydroxyalkanoates are a class of microbial polyesters synthesized by some bacteria as intracellular storage compounds. PHAs are composed of various hydroxyalkanoic acids and can serve as markers for the presence of bacteria capable of producing these polymers [79-82].

13. Omega-9 fatty acids: Omega-9 fatty acids are monounsaturated fatty acids with a double bond at the ninth carbon atom from the omega end of the carbon chain. Although omega-9 fatty acids are also found in higher organisms, they can be used as markers for certain microbial species or strains [83-86].

14. Conjugated fatty acids: Conjugated fatty acids are characterized by the presence of several conjugated double bonds. These fatty acids are produced by some bacteria and can serve as markers for certain microbial groups. Examples include conjugated linoleic acid (CLA) produced by lactic acid bacteria [87,88].

15.Iso/ anteiso fatty acids: Iso and anteiso fatty acids are branched chain fatty acids that contain a methyl group attached to a carbon atom adjacent to a carboxyl group. These fatty acids are commonly found in bacteria and can serve as markers for specific bacterial groups or taxonomic identification. Examples include iso-C15:0 and iso-C17:0 [89-90].

16.3-hydroxy fatty acids: 3-hydroxy fatty acids are fatty acids containing a hydroxyl group at the third carbon atom. These fatty acids are produced by some bacteria as components of complex lipids and can act as markers for certain species or groups of microbes. Examples include 3-hydroxydecanoic acid (C10:0 3-OH) and 3-hydroxytetradecanoic acid (C14:0 3-OH) [49,75,91-93].

17. Mycolic acids: Mycolic acids are long chain fatty acids with a unique structure that are found in the cell walls of some bacteria, especially mycobacteria. These fatty acids contribute to the characteristic waxy appearance of the cell wall and can serve as markers for the presence of mycobacteria [94,95].

18. Gopanoids: hopanoids are bacterial lipids that are structurally similar to sterols that are found in eukaryotes. These lipids are widely distributed in bacteria and can act as biomarkers for bacterial populations. Hopanoids have been used as markers for the presence of certain groups of bacteria, such as cyanobacteria and some proteobacteria [96-99].

19. Wax esters: Wax esters are ester compounds formed by the esterification of fatty acids with long chain alcohols. Some bacteria and fungi can produce wax esters, and their presence can serve as a marker for the activity of certain micro-organisms or metabolic processes. Wax esters have been studied in various environments, including marine systems and soil microbial communities [13,100-104].

20. Sterols: Although sterols are more commonly associated with eukaryotes, some bacteria, such as some *Mycoplasma* species, are able to synthesize sterol-like compounds called bacteriohopan polyols (BHPs). These compounds can serve as markers for the presence of certain groups of bacteria and have been used in paleoecological studies as biomarkers of microbial activity [15,105-108].

21. Phospholipids of fatty acids (PLFA): Phospholipids of fatty acids are the fatty acid components of phospholipids, which are the main constituents of the cell membranes of microorganisms. PLFA analysis can provide insight into microbial biomass, community structure, and functional diversity. Various PLFAs can be used as markers for certain microbial groups or metabolic activity [39,46,109-115].

22. Lipopolysaccharides (LPS): Lipopolysaccharides are complex molecules found in the outer membrane of Gram-negative bacteria. They consist of a lipid component (lipid A) and a polysaccharide component. The fatty acid composition of lipid A can vary between bacterial species and strains, and LPS analysis can provide markers for the presence and identification of specific Gram-negative bacteria [116-119].

23. Ether lipids: While most microorganisms contain ester-linked fatty acids, some archaea, such as methanogens, produce ether lipids in their cell membranes. Ether lipids have unique structural characteristics and can serve as markers for the presence of specific archaeal groups [51,120].

24. Very Long Chain Fatty Acids (VLCFA): Very long chain fatty acids generally refer to fatty acids with a carbon chain length of 20 or more carbon atoms. These fatty acids are found in various microorganisms and can serve as markers for specific microbial groups or metabolic activity. VLCFAs have been studied in the context of soil microbial communities and their potential role in nutrient cycling [64,69,121-124].

25. Diacylglycerols: Diacylglycerols are lipid molecules composed of two fatty acid chains attached to a glycerol backbone. They are important intermediates in lipid metabolism and membrane biosynthesis [13,121,125-127].

Advantages of GC-MS [1-4] :

- **High Sensitivity:** GC-MS is one of the most sensitive methods for analyzing organic compounds. It allows the detection and determination of substances at very low concentrations, which is especially important for medical, food and environmental analysis.

- **High specificity:** GC-MS provides excellent recognition specificity and a unique ability to determine the molecular structure of the analyzed compounds. This allows you to get more accurate and reliable results.

- **Ability to analyze a wide range of compounds:** GC-MS can be used to analyze various classes of organic compounds, including volatile and thermostable substances, aromatic and heterocyclic compounds, polymers, and even living tissues.

- **Fast and automated:** GC-MS is a fast and relatively easy-to-use method of analysis. Modern instruments have the ability to automate sample insertion and data processing, minimizing human error and increasing productivity.

Limitations of GC/MS [5-9] :

- **High equipment cost:** Acquisition and maintenance of GC/MS equipment is a significant financial investment, especially for high performance and multi-channel systems.

➤ **Difficulty in interpreting data:** Interpreting mass spectral data can be complex and requires specific knowledge and skills. In some cases, additional analysis or database comparison may be required to fully identify compounds.

➤ **Insufficient Resolution:** Some compounds may experience peak overlapping problems, especially when analyzing complex mixtures. This can complicate identification and quantification.

➤ **Possibility of analyte degradation:** High temperatures, use of catalytic surfaces, and other operating conditions in GC/MS can degrade some compounds. This may lead to false results or the inability to analyze certain substances.

Future prospects for GC-MS and conclusions.

The future prospects of gas chromatography -mass spectrometry (GC-MS) promise to be very promising in many scientific and industrial fields. Here are a few key takeaways about the future of this technology:

➤ **Instrumentation Developments:** Significant developments in GC/MS equipment are expected over time, including improved detector sensitivity, increased resolution, and increased analysis speed. This will allow researchers to obtain more accurate and reliable analysis results.

➤ **Expanding Applications:** GC/MS has already been successfully applied in a variety of applications including food, environmental, pharmaceutical, oil refining, and more. In the future, GC-MS is expected to find new applications such as forensics, medicine, and biological research.

➤ **Methodological evolution:** With the development of GC/MS, new and improved methods of analysis are expected. For example, multivariate GC/MS allows you to get more detailed information about the composition of samples, as well as integrate data from various sources for a more complete analysis.

➤ **Application of artificial intelligence and machine learning:** In the future, the use of artificial intelligence and machine learning methods in the analysis of GC-MS data is expected to increase. This will automate and speed up the analysis process, as well as facilitate the interpretation of the results.

All in all, the future prospects of GC/MS promise to be very attractive and useful for research and industry. This technology will continue to evolve, opening up new possibilities for analysis and allowing a more complete understanding of complex chemical and biological systems.

The choice and identification of specific fatty acid markers depends on the focus of the study and the target microorganisms. The use of fatty acids as markers in microbial analysis provides valuable insight into microbial community structure, metabolic activity, and ecological functions. Advanced analytical techniques such as chromatography, mass spectrometry, and lipidomics have contributed greatly to the identification and characterization of fatty acid markers in microorganisms.

Overall, the analysis of fatty acids as markers in microorganisms provides valuable information for microbial identification, analysis of community structure, and understanding of the functional roles of microorganisms in various environments. Techniques such as gas chromatography-mass spectrometry (GC-MS) are commonly used to analyze fatty acids, allowing researchers to gain insight into microbial communities and their role in plant-microbial interactions, environmental processes, and human health.

Literature

1. Tsavkelova E. L. et al. Microorganisms-producers of plant growth stimulants and their practical application (review) // Applied Biochemistry and Microbiology. - 2006. - T. 42. - No. 2. - S. 133-143.

2. Maksimov I. V., Abizgildina R. R., Pusenkova L. I. Microorganisms that stimulate plant growth as an alternative to chemical means of protection against pathogens (review) // *Applied Biochemistry and Microbiology*. - 2011. - T. 47. - No. 4. - S. 373-385.
3. Levitin M. M. Microorganisms in the conditions of global climate change // *Agricultural biology*. – 2015. – no. 5. - S. 641-647.
4. Zaikin VG Chromato-mass spectrometry in Russia // *Journal of Analytical Chemistry*. - 2011. - T. 66. - No. 11. - S. 1205-1209.
5. Yashin Ya. I., Yashin A. Ya. Analytical chromatography. Methods, equipment, application // *Advances in Chemistry*. - 2006. - T. 75. - No. 4. - S. 366-379.
6. Gladilovich V. D., Podolskaya E. P. Possibilities of using the GC-MS method (Review) // *Scientific Instrumentation*. - 2010. - T. 20. - No. 4. - S. 36-49.
7. Hübschmann HJ Handbook of GC-MS: fundamentals and applications. – John Wiley & Sons, 2015.
8. Pisanov R. V. et al. Identification of microorganisms using gas chromatography-mass spectrometry // *Journal of Microbiology, Epidemiology and Immunobiology*. – 2020. – no. 4. - S. 356-362.
9. Savelyeva E. I., Gavrilova O. P., Gagkaeva T. Yu. Application of solid-phase microextraction in combination with gas chromatography-mass spectrometry for the study of volatile biosynthesis products released by plants and microorganisms // *Journal of Analytical Chemistry*. - 2014. - T. 69. - No. 7. - S. 675-675.
10. Veselova M. A., Plyuta V. A., Khmel I. A. Volatile substances of bacteria: structure, biosynthesis, biological activity // *Microbiology*. - 2019. - T. 88. - No. 3. - S. 272-287. zzzz
11. Luca A., Kjør A., Edelenbos M. Volatile organic compounds as markers of quality changes during the storage of wild rocket // *Food Chemistry*. - 2017. - V. 232. - P. 579-586.
12. Barac T. et al. Engineered endophytic bacteria improve phytoremediation of water-soluble, volatile, organic pollutants // *Nature biotechnology*. – 2004. – V. 22. – no. 5.-P. __ 583-588.
13. Rozentsvet O. A., Fedoseeva E. V., Terekhova V. A. Lipid biomarkers in the ecological assessment of soil biota : analysis of fatty acids // *Successes of modern biology*. - 2019. - T. 139. - No. 2. - S. 161-177.
14. Makhutova O. N., Sushchik N. N., Kalacheva G. S. Informativity of the composition of fatty acids of triacylglycerols and polar lipids of seston in the analysis of the feeding spectrum of microzooplankton of the small Bugach reservoir // *Reports of the Academy of Sciences. - Federal State Budgetary Institution "Russian Academy of Sciences"*, 2004. - T. 395. - No. 4. - S. 562-565.
15. Verkhovtseva NV, Osipov GA Method of gas chromatography-mass spectrometry in the study of microbial communities of soils of agrocenosis // *Problems of agrochemistry and ecology*. - 2008. - no. 1. - S. 51-54.
16. Pollierer MM, Scheu S., Haubert D. Taking it to the next level: trophic transfer of marker fatty acids from basal resource to predators // *Soil Biology and Biochemistry*. - 2010. - T. 42. - no. 6. - S. 919-925.
17. Loshchinina E. A., Nikitina V. E. Changes in the content of stress metabolites of xylophilic basidiomycetes under the influence of microorganisms // *Mechanisms of resistance of plants and microorganisms to unfavorable environmental*. - 2018. - S. 479.
18. Chetverikov S.P. New metabolites of bacteria of the genus *Pseudomonas* - inhibitors of the growth of phytopathogenic fungi : dis . - Ufa : [Inst. of biochemistry and genetics Ufim . scientific Center of the Russian Academy of Sciences], 2004.

19. Iturriaga G., Suárez R., Nova-Franco B. Trehalose metabolism: from osmoprotection to signaling // *International journal of molecular sciences*. - 2009. - T . 10. - no. 9. - S. 3793-3810.
20. Hennion N. et al. Sugars en route to the roots. Transport, metabolism and storage within plant roots and towards microorganisms of the rhizosphere // *Physiologia plantarum* . - 2019. - T . 165. - no. 1. - S. 44-57.
21. Haggag WM et al. Biotechnological aspects of microorganisms used in plant biological control // *American-Eurasian Journal of Sustainable Agriculture*. - 2007. - T . 1. - no. 1. - S. 7-12.
22. Smirnova O. G., Kochetov A. V. Plant cell wall and mechanisms of resistance to pathogens // *Vavilov Journal of Genetics and Breeding*. - 2016. - T. 19. - No. 6. - S. 715-723.
23. Kim D. et al. Glycopolymers cell wall - a chemotaxonomic marker of actinobacteria of the genus *Clavibacter* // VII Pushchino Conference "Biochemistry, Physiology and Biospheric Role of Microorganisms". School-conference for young scientists, graduate students and students "Genetic technologies in microbiology and microbial diversity". - 2021. - S. 50-51.
24. Bacete L. et al. Arabidopsis response regulator 6 (ARR6) modulates plant cell-wall composition and disease resistance // *Molecular Plant-Microbe Interactions*. - 2020. - T . 33. - no. 5. - S. 767-780.
25. Vargas- Asensio G. et al. Uncovering the cultivable microbial diversity of costa rican beetles and its ability to break down plant cell wall components // *PLoS One*. - 2014. - T . 9. - no. 11. - P. e113303.
26. Solovieva IV et al. Biological properties of lactobacilli . Prospects for use in the laboratories of Rospotrebnadzor express methods of amplification of nucleic acids (NAAT) in the quality control of food products, dietary supplements for food, dosage forms containing lactobacilli // *Journal of MediAl* . – 2014. – no. 2 (12). - S. 29-44.
27. Plotnikov VK Nanobiotechnological methods for the study of nucleic acids and prospects for their practical application // *News of the Timiryazev Agricultural Academy*. - 2009. - no. 4. - S. 58-70.
28. Malik AA et al. Rhizosphere bacterial carbon turnover is higher in nucleic acids than membrane lipids: implications for understanding soil carbon cycling // *Frontiers in Microbiology*. - 2015. - T . 6. - S. 268.
29. Vlassov VV, Laktionov PP, Rykova EY Extracellular nucleic acids // *Bioessays* . - 2007. - T . 29. - no. 7. - S. 654-667.
30. Vladimirova AS et al. Fluorescent proteins as markers of conjugative interaction of nodule bacteria // *Experimental plant biology: fundamental and applied aspects*. - 2017. - S. 127-127.
31. Maksimov IV et al. Plant growth-stimulating bacteria in the regulation of plant resistance to stress factors // *Plant Physiology*. - 2015. - T. 62. - No. 6. - S. 763-775.
32. Chesnokov Yu. V. Biochemical markers in genetic studies of cultivated plants: applicability and limitations // *Agricultural biology*. - 2019. - T. 54. - No. 5. - S. 863-874.
33. Jain A. et al. Microbial consortium-induced changes in oxidative stress markers in pea plants challenged with *Sclerotinia sclerotiorum* // *Journal of Plant Growth Regulation*. - 2013. - T . 32. - S. 388-398 .
34. Olempska -Beer ZS et al. Food-processing enzymes from recombinant microorganisms—a review // *Regulatory toxicology and Pharmacology*. - 2006. - T . 45. - no. 2. - S. 144-158.
35. Khoseeva E. V., Zimina Yu. A., Sroslova G. A. Oxidative stress of plants: chemistry, physiology, methods of protection // *Natural systems and resources*. - 2020. - T. 10. - No. 4. - S. 30-43.

36. Malinchik M. A. et al. Application of the 16S rRNA RFLP gene analysis method for the identification of bacteria of the genus *Rhizobium* : dis . – Siberian Federal University, 2017.
37. Pospíšil P., Prasad A., Rác M. Mechanism of the formation of electronically excited species by oxidative metabolic processes: role of reactive oxygen species //Biomolecules. - 2019. - T . 9. - no. 7. - S. 258.
38. Wilson SA, Roberts SC Recent advances towards development and commercialization of plant cell culture processes for the synthesis of biomolecules // Plant biotechnology journal. - 2012. - T . 10. - no. 3. - S. 249-268.
39. Shipko E. S., Duvanova O. V. Changing the spectrum of fatty acids as one of the mechanisms of adaptation / persistence of microorganisms // Journal of Microbiology, Epidemiology and Immunobiology. – 2019. – no. 5. - S. 109-118.
40. Zelles L. Phospholipid fatty acid profiles in selected members of soil microbial communities // Chemosphere. - 1997. - T . 35. - no. 1-2. - S. _ 275-294.
41. Bessa RJB et al. Using microbial fatty acids to improve understanding of the contribution of solid associated bacteria to microbial mass in the rumen // Animal Feed Science and Technology. - 2009. - T . 150. - no. 3-4. - S. _ 197-206.
42. Vlaeminck B. et al. Factors affecting odd-and branched-chain fatty acids in milk: A review //Animal feed science and technology. - 2006. - T . 131. - no. 3-4. - S. _ 389-417.
43. Trefflich I. et al. Short-and branched-chain fatty acids as fecal markers for microbiota activity in vegans and omnivores //Nutrients. - 2021. - T . 13. - no. 6. - S. 1808.
44. Choi BSY et al. Feeding diversified protein sources exacerbates hepatic insulin resistance via increased gut microbial branched-chain fatty acids and mTORC1 signaling in obese mice //Nature communications. - 2021. - T . 12. - no. 1. - S. 3377.
45. Zakharchenko N. S. et al. Obtaining biosafe marker-free *Camelina sativa* with increased resistance to phytopathogens // Mechanisms of resistance of plants and microorganisms to unfavorable environmental . - 2018. - S. 350.
46. Hinojosa MB et al. Microbial Response to Heavy Metal–Polluted Soils: Community Analysis from Phospholipid - Linked Fatty Acids and Ester - Linked Fatty Acids Extracts // Journal of Environmental Quality. - 2005. - T . 34. - no. 5. - S. 1789-1800.
47. Unger IM, Kennedy AC, Muzika RM Flooding effects on soil microbial communities // Applied Soil Ecology. - 2009. - T . 42. - no. 1. - S. 1-8.
48. Sakharuta I. Yu., Lagodich O. V. The use of molecular markers for the study of ISR // Biotechnology in crop production, animal husbandry and agricultural microbiology. - 2019. - S. 214-215.
49. Liu-Lyanmin E. I. Identification and determination of lipid components of hilly permafrost peatlands // Actual problems of biology and ecology. - 2021. - S. 78-81.
50. Saini RK et al. Omega- 3 polyunsaturated fatty acids (PUFAs): Emerging plant and microbial sources, oxidative stability, bioavailability, and health benefits— A review //Antioxidants. - 2021. - T . 10. - no. 10. - S. 1627.
51. Rizzo G., Baroni L., Lombardo M. Promising sources of plant-derived polyunsaturated fatty acids: A narrative review // International Journal of Environmental Research and Public Health. - 2023. - T . 20. - no. 3. - S. 1683.
52. Sushchik N. N. The role of essential fatty acids in trophometabolic interactions in freshwater ecosystems (review) //Journal of General Biology. - 2008. - T. 69. - No. 4. - S. 299-316.
53. Caligiani A., Lolli V. Cyclic fatty acids in food: An under investigate class of fatty acids //Biochemistry and Health Benefits of Fatty Acids. - 2018. - T . 2018.

54. Munoz-Rojas J. et al. Involvement of cyclopropane fatty acids in the response of *Pseudomonas putida* KT2440 to freeze-drying // *Applied and Environmental Microbiology*. - 2006. - T . 72. - no. 1. - S. 472-477.
55. Dahlquist A. et al. A new class of enzymes in the biosynthetic pathway for the production of triacylglycerol and recombinant DNA molecules encoding these enzymes. – 2006. (Patent)
56. Tyagi P. et al. Hydroxy fatty acids in snow pit samples from Mount Tateyama in central Japan: Implications for atmospheric transport of microorganisms and plant waxes associated with Asian dust // *Journal of Geophysical Research: Atmospheres*. - 2016. - T . 121. - no. 22. - S. 13.641-13.660.
57. Tyagi P. et al. Impact of biomass burning on soil microorganisms and plant metabolites: A view from molecular distributions of atmospheric hydroxy fatty acids over Mount Tai // *Journal of Geophysical Research: Biogeosciences* . - 2016. - T . 121. - no. 10. - S. 2684-2699.
58. Bikkina P. et al. Decadal Variations in Hydroxy Fatty Acids Over Chichijima Island in the North Pacific: Long - Term Seasonal Variability in Plant and Microbial Markers // *Journal of Geophysical Research: Atmospheres*. - 2021. - T . 126. - no. 21. - P. e2020JD033347.
59. Glyzina O. Yu. et al. Changes in the lipid composition of freshwater sponges with an increase in environmental temperature // *Ecology*. – 2016. – no. 2. - S. 152-155.
60. Silina A. V., Zhukova N. V. Benthic association of a bivalve mollusk with a borer polychaete and their potential food sources // *Oceanology*. - 2012. - T. 52. - No. 5. - S. 700-700.
61. Kirichenko K. A. et al. Comparative analysis of the fatty acid composition of coastal water *Typha latifolia* , submerged by *Ceratophyllum demersum* and water form *Veronica anagallis-aquatica* of water bodies of the Baikal region // *Chemistry of plant raw materials*. – 2019. – no. 4. - S. 119-128.
62. Kim EJ et al. Fatty acid profiles associated with microbial colonization of freshly ingested grass and rumen biohydrogenation // *Journal of Dairy Science*. - 2005. - T . 88. - no. 9. - S. 3220-3230.
63. Zhang LS et al. Microbial synthesis of functional odd-chain fatty acids: a review // *World Journal of Microbiology and Biotechnology*. - 2020. - T . 36. - S. 1-9.
64. Řezanka T., Sigler K. Odd-numbered very-long-chain fatty acids from the microbial, animal and plant kingdoms // *Progress in lipid research*. - 2009. - T . 48. - no. 3-4. – S. 206-238.
65. Sushchik N. N. The role of essential fatty acids in trophometabolic interactions in freshwater ecosystems (review) // *Journal of General Biology*. - 2008. - T. 69. - No. 4. - S. 299-316.
66. Fan Y. et al. Week-old chicks with high *Bacteroides* abundance have increased short-chain fatty acids and reduced markers of gut inflammation // *Microbiology Spectrum*. - 2023. - T . 11. - no. 2. - S. e03616-22.
67. Nagpal R. et al. Modified Mediterranean- ketogenic diet modulates gut microbiome and short-chain fatty acids in association with Alzheimer's disease markers in subjects with mild cognitive impairment // *EBioMedicine* . - 2019. - T . 47. - S. 529-542 .
68. Zhang YM et al. Relationships between rumen microbes, short-chain fatty acids, and markers of white adipose tissue browning during the cold season in grazing Mongolian sheep (*Ovis aries*) // *Journal of Thermal Biology*. - 2022. - T . 110. - S. 103386.
69. Kormilets O.N. Fatty acids in food webs of inland water ecosystems : dis . – Siberian Federal University, 2019.
70. Makhutova O. N., Gladyshev M. I. Essential polyunsaturated fatty acids in the physiology and metabolism of fish and humans: significance, needs, sources // *Russian Journal of Physiology*. IM Sechenov. - 2020. - T. 106. - No. 5. - S. 601-621-601-621.

71. Krivova Z. V., Maltsev E. I., Kulikovskiy M. S. Comparison of fatty acid profiles of different strains of the diatom *Cyclotella meneghiniana* Kützinger from the salt lake Takhilt-Nuur (Mongolia) // Problems of Botany of Southern Siberia and Mongolia. - 2021. - T. 20. - No. 1. - S. 246-248.
72. Adarme -Vega TC et al. Microalgal biofactories : a promising approach towards sustainable omega-3 fatty acid production //Microbial cell factories. - 2012. - T . 11. - no. 1. - S. 1-10.
73. Lee JH et al. Omega-3 fatty acids: cardiovascular benefits, sources and sustainability //Nature Reviews Cardiology. - 2009. - T . 6. - no. 12. - S. 753-758.
74. Ye VM, Bhatia SK Metabolic engineering for the production of clinically important molecules: Omega - 3 fatty acids, artemisinin , and taxol //Biotechnology journal. - 2012. - T . 7. - no. 1. - S. 20-33.
75. Gessler N. N. et al. Oxylipins and ways of their synthesis in fungi //Applied Biochemistry and Microbiology. - 2017. - T. 53. - No. 6. - S. 568-579.
76. Starodumova I. P. Development of a classification system for actinobacteria of the genus *Rathayibacter* : dis . - M.: Federal Research Center "Fundamental Foundations of Biotechnology" of the Russian Academy of Sciences, Institute of Microbiology. SN Vinogradsky, 2018.
77. Varbanets L. D., Vasiliev V. N., Brovarskaya O. S. Characterization of lipopolysaccharides *Ralstonia solanacearum* //Microbiology. - 2003. - T. 72. - No. 1. - S. 19-25.
78. Ibekwe AM, Kennedy AC Fatty acid methyl ester (FAME) profiles as a tool to investigate community structure of two agricultural soils //Plant and Soil. - 1999. - T . 206. - S. 151-161 .
79. Volova T. G. Modern biomaterials: world trends, place and role of microbial polyhydroxyalkanoates // Journal of the Siberian Federal University. Biology. - 2014. - T. 7. - No. 2. - S. 103-133.
80. Parshina V. V., Dyatlova Yu. A., Tugarova A. V. Ikfurier spectroscopic analysis of the accumulation of poly-3-hydroxybutyrate by *Azospirillum* cells *brasiliense* with different duration of cultivation and ammonium concentration in the nutrient medium // Bulletin of the Saratov University. New episode. Series Chemistry. Biology. Ecology. - 2018. - T. 18. - No. 3. - S. 331-335.
81. Valentin HE et al. PHA production, from bacteria to plants //International Journal of Biological Macromolecules. - 1999. - T . 25. - no. 1-3. - S. _ 303-306.
82. Gasser I., Müller H., Berg G. Ecology and characterization of polyhydroxyalkanoate - producing microorganisms on and in plants //FEMS Microbiology Ecology. - 2009. - T . 70. - no. 1. - S. 142-150.
83. Walsh , T. A. et al. Production of DHA and other LC-PUFAs in plants. – 2018. (Ratent)
84. Farag MA, Gad MZ Omega-9 fatty acids: Potential roles in inflammation and cancer management // Journal of Genetic Engineering and Biotechnology. - 2022. - T . 20. - no. 1. - S. 1-11.
85. HaghighiTM et al. Adaptation of *Glycyrrhiza glabra* L. to water deficiency based on carbohydrate and fatty acid quantity and quality //Scientific Reports. - 2023. - T . 13. - no. 1. - S. 1766.
86. Wang K. et al. Engineering the lipid and fatty acid metabolism in *Yarrowia lipolytica* for sustainable production of high oleic oils //ACS Synthetic Biology. - 2022. - T . 11. - no. 4. - S. 1542-1554.

87. Yakimova T. V., Nasanova O. N., Vengerovsky A. I. Antidiabetic effect of some medicinal plants (based on publications of the last 15 years) // Plant Resources. - 2016. - T. 52. - No. 1. - S. 3-19.
88. Dewhurst RJ et al. Forage breeding and management to increase the beneficial fatty acid content of ruminant products //Proceedings of the Nutrition society. - 2003. - T . 62. - no. 2. - S. 329-336.
89. Rodkina S.A. Fatty acids and other lipids of sea sponges //Biology of the sea. - 2005. - T. 31. - No. 6. - S. 387-397.
90. Kaneda T. Iso -and anteiso -fatty acids in bacteria: biosynthesis, function, and taxonomic significance //Microbiological reviews. - 1991. - T . 55. - no. 2. - S. 288-302.
91. Motelskaya V. A. et al. Modern methods of laboratory diagnosis of chlamydia // Journal of Microbiology, Epidemiology and Immunobiology. - 2008. - no. 4. - S. 111-117.
92. Jenske R., Vetter W. Enantioselective analysis of 2-and 3-hydroxy fatty acids in food samples // Journal of agricultural and food chemistry. - 2008. - T . 56. - no. 24. - S. 11578-11583.
93. Goossens H. et al. Lipids and their mode of occurrence in bacteria and sediments—II. Lipids in the sediment of a stratified, freshwater lake //Organic Geochemistry. - 1989. - T . 14. - no. 1. - S. 27-41.
94. Agafonova N. V. Taxonomic and functional characteristics of aerobic methylotrophic bacteria-phytosymbionts : dis . - - Pushchino: IBFM RAS, 2017. - 156 s, 2017.
95. Stainsby FM et al. Dispelling the “ Nocardia amarae ” myth: a phylogenetic and phenotypic study of mycolic acid-containing actinomycetes isolated from activated sludge foam // Water Science and Technology. - 2002. - T . 46. - no. 1-2. – S. 81-90.
96. Duchko M.A. Geochemistry of biomarkers in peats of the southeastern part of Western Siberia: dissertation for the degree of candidate of geological and mineralogical sciences: spec. 25.00. 09 : dis . – 2016.
97. Karpunina L. V. Exopolysaccharides of bacteria of the genera Xanthobacter and Ancylobacter : Characteristics and their biological properties: dissertation . for the competition of a candidate of biological sciences: spec. 03.02.03. - Saratov, 2019. - 121 p.
98. Plemenkov VV, Tevs OA Medico-biological properties and prospects of terpenoids (isoprenoids) // Chemistry of vegetable raw materials. – 2014. – no. 4. - S. 5-20.
99. Belin BJ et al. Hopanoid lipids: from membranes to plant–bacteria interactions //Nature Reviews Microbiology. - 2018. - T . 16. - no. 5. - S. 304-315.
100. Botirov E. Kh., Bonacheva V. M., Kolomiets N. E. Chemical composition and biological activity of plant metabolites of the genus Equisetum L // Chemistry of plant raw materials. – 2021. – no. 1. - S. 5-26.
101. Alotaibi SS et al. Transcriptome analysis of jojoba (Simmondsia chinensis) during seed development and liquid wax ester biosynthesis //Plants. - 2020. - T . 9. - no. 5. - S. 588.
102. Simoneit BRT Organic matter of the troposphere—V: application of molecular marker analysis to biogenic emissions into the troposphere for source reconciliations //Journal of Atmospheric Chemistry. - 1989. - T . 8. - S. 251-275 .
103. Saliot A. Sources markers in aerosols, oceanic particles and sediments //EPJ Web of Conferences. - EDP Sciences, 2009. - T . 1. - S. 189-197 .
104. Mayes RW et al. Discrimination of domestic garden soils using plant wax compounds as markers //Criminal and environmental soil forensics. - 2009. - S. 463-476 .

105. Osipov GA A method of calibrating a gas chromatography-mass spectrometry (GC-MS) system equipped with special software for determining microbial markers in a test sample of a material of biological origin. – 2013 (Ratent).
106. Reichel R. et al. Effects of slurry from sulfadiazine-(SDZ) and difloxacin -(DIF) medicated pigs on the structural diversity of microorganisms in bulk and rhizosphere soil //Soil Biology and Biochemistry. - 2013. - T . 62. - S. 82-91 .
107. Ourisson G., Rohmer M., Poralla K. Prokaryotic hopanoids and other polyterpenoid sterol surrogates // Annual Reviews in Microbiology. - 1987. - T . 41. - no. 1. - S. 301-333.
108. S'amaj J. et al. Endocytosis, actin cytoskeleton, and signaling // Plant physiology. - 2004. - T . 135. - no. 3. - S. 1150-1161.
109. Akhmetshina E. A., Sirotkin A. S. Analysis of phospholipid fatty acids of microorganisms as environmental biomarkers // Bulletin of the Kazan Technological University. - 2014. - T. 17. - No. 19. - S. 233-236.
110. Evdokimov I. V., Larionova A. A., Stulin A. F. Turnover of "new" and "old" carbon in the biomass of soil microorganisms // Microbiology. - 2013. - T. 82. - No. 4. - S. 489-489.
111. Evdokimov I. V. Methods for determining the biomass of soil microorganisms // Russian Journal of Ecosystem Ecology . – 2018. – no. 3. - S. 1-20.
112. Machekhina VV Determination of the composition of fatty acids in sapropel by chromato-mass spectrometry using various extraction methods: master's thesis in the field of study: 04.04. 01-Chemistry. – 2016.
113. Willers C., Jansen van Rensburg PJ, Claassens S. Phospholipid fatty acid profiling of microbial communities—a review of interpretations and recent applications //Journal of applied microbiology. - 2015. - T . 119. - no. 5. - S. 1207-1218.
114. Miura T. et al. Comparison of fatty acid methyl ester methods for characterization of microbial communities in forest and arable soil: Phospholipid fraction (PLFA) versus total ester linked fatty acids (EL-FAME) // Pedobiologia . - 2017. - T . 63. - S. 14-18 .
115. Calderon FJ et al. Short - term dynamics of nitrogen, microbial activity, and phospholipid fatty acids after tillage //Soil Science Society of America Journal. - 2001. - T . 65. - no. 1. - S. 118-126.
116. Ignatov V. V. et al. Characteristics of the composition of fatty acids of lipids A of lipopolysaccharides of bacteria of the genus Azospirillum // Bulletin of the Saratov University. New episode. Series Chemistry. Biology. Ecology. - 2009. - T. 9. - No. 1. - S. 36-41.
117. Sigida E. N. et al. Comparative characteristics of lipopolysaccharides of bacteria of the Azospirillum strain brasiliense Sp7 and its spontaneous mutant Sp7. K2 //News of the Saratov University. New episode. Series Chemistry. Biology. Ecology. - 2012. - T. 12. - No. 1. - S. 61-65.
118. Lagares A. et al. A Rhizobium meliloti lipopolysaccharide mutant altered in competitiveness for nodulation of alfalfa // Journal of bacteriology. - 1992. - T . 174. - no. 18. - S. 5941-5952.
119. Ramos Solano B. et al. Systemic disease protection elicited by plant growth promoting rhizobacteria strains: relationship between metabolic responses, systemic disease protection, and biotic elicitors //Phytopathology. - 2008. - T . 98. - no. 4. - S. 451-457.
120. Meziane T., Tsuchiya M. Fatty acids as tracers of organic matter in the sediment and food web of a mangrove/intertidal flat ecosystem, Okinawa, Japan // Marine Ecology Progress Series. - 2000. - T. 200. - S. 49-57 .

121. Makhutova O. N., Pryanichnikova E. G., Lebedeva I. M. Comparison of feeding spectra of zebra mussel *Dreissena polymorpha* and *Dreissena bugensis* by biochemical markers // Siberian Ecological Journal. - 2012. - T. 19. - No. 4. - S. 619-631.
122. Kyselová L., Vítová M., Řezanka T. Very long chain fatty acids // Progress in Lipid Research. - 2022. - S. 101180.
123. Volkman JK et al. Microbial lipids of an intertidal sediment—I. Fatty acids and hydrocarbons // Geochimica et cosmochimica acta . - 1980. - T . 44. - no. 8. - S. 1133-1143.
124. Hatamoto M. et al. Diversity of anaerobic microorganisms involved in long-chain fatty acid degradation in methanogenic sludges as revealed by RNA-based stable isotope probing // Applied and environmental microbiology. - 2007. - T . 73. - no. 13. - S. 4119-4127.
125. Makhutova O. N. et al. Seasonal dynamics of the nutritional spectrum of *Dreissena polymorpha* in the Rybinsk Reservoir // Reports of the Academy of Sciences. - Federal State Budgetary Institution "Russian Academy of Sciences", 2008. - T. 423. - No. 5. - S. 710-713.
126. Kim MJ et al. Gene silencing of Sugar-dependent 1 (JcSDP1), encoding a patatin -domain triacylglycerol lipase, enhances seed oil accumulations in *Jatropha curcas* // Biotechnology for biofuels. - 2014. - T . 7. - S. 1-16.
127. LuCL et al. Expression pattern of diacylglycerol acyltransferase-1, an enzyme involved in triacylglycerol biosynthesis, in *Arabidopsis thaliana* // Plant molecular biology. - 2003. - T . 52. - S. 31-41 .

CENTRAL ASIAN
STUDIES