



Impact of Vitamin D Deficiency on Pregnancy

1. Sharipova N. M.

Received 2nd Aug 2023,
Accepted 19th Sep 2023,
Online 19th Oct 2023

¹ Bukhara State Medical Institute, Bukhara,
Uzbekistan

Abstract: Relevance. Recently, much attention has been paid to studying the prevalence of vitamin D insufficiency and deficiency in the population. The steps involved in the biological effects of vitamin D are being studied extensively. It should be recognized that this disorder is of interest to both obstetrician-gynecologists and other specialists, since vitamin D deficiency is associated with oncological, neurological, cardiovascular and autoimmune diseases, as well as various pathologies in the field of obstetrics and gynecology. Recently, there has been a sharp increase in interest in the "hormone D" - calcitriol and its receptors (VDR) as risk factors for the development of a number of obstetric complications. Vitamin D receptors have been found in the ovaries, uterus, placenta, and pituitary gland. According to some studies, vitamin D deficiency during pregnancy increases the risk of preterm birth. It is involved in the processes of angiogenesis, inhibition of cell proliferation, maintains genetic homeostasis and final differentiation of fetuses, affects the production of macrophages, the functioning of the pancreas, and the reninangiotensin system (1, 16,31).

Key words: vitamin D deficiency, 25-hydroxyvitamin D, general risk groups, pregnancy, preterm birth.

The incidence of vitamin D deficiency in the population varies from 20% to 90% (2,30). The prevalence of vitamin D deficiency and insufficiency during pregnancy ranges from 19.0 to 96.0% in Europe, from 27.0 to 91.0% in the United States, from 39.0 to 65.0% in Canada, in Australia and New Zealand - from 25.0 to 87.0% and in Asia - from 45.0 to 100.0% (5). According to a meta-analysis of 195 vitamin D status studies conducted in 44 countries and involving more than 168,000 people, the average serum level of 25-hydroxyvitamin D (25(OH)D) in the population varies widely from 1.9 to 54.6 ng/ml. At the same time, 37.3% of researchers reported vitamin D deficiency (<20 ng/ml) in the population. It has been shown that vitamin D deficiency in the body of a pregnant woman is associated with an increased risk of premature birth, the development of placental insufficiency, preeclampsia, gestational diabetes mellitus, bacterial vaginosis, impaired contractile activity of the uterus, and an increase in the frequency of cesarean sections. In the fetus and newborn it leads to low birth weight, decreased femoral growth in utero, infant heart failure, craniotabes, acute lower respiratory tract

infection, and hypocalcemia. Vitamin D deficiency is considered to be a negative factor affecting women's reproductive health (7,11,29).

The review is devoted to calcitriol-dependent mechanisms of the pathogenesis of such obstetric complications as spontaneous miscarriage, premature birth, fetal growth restriction, preeclampsia, perinatal infections and gestational diabetes mellitus. In light of the achievements of recent years, the role of vitamin D in the regulation of reproductive function has been proven. As a steroid hormone, this vitamin is necessary for a wide range of physiological processes. Vitamin D deficiency is currently a worldwide health problem and causes a very large percentage of acute and chronic pathologies. It actively influences various metabolic processes and takes part in the regulation of cell growth and functioning in the human body (6,32). It was reported that pregnant women with serum vitamin D concentrations less than 20 ng/mL were 3.81 times more likely to have preterm birth than women with vitamin D concentrations greater than 40 ng/mL (18,28,33).

Vitamin D is formed in the skin under the influence of ultraviolet radiation or comes from food, then a chain of metabolic processes occurs with the formation of active metabolites of vitamin D, which, together with parathyroid hormone and calcitonin, provide regulation of calcium and phosphate metabolism - the so-called classical action of vitamin D. In pregnant women, syncytiotrophoblast is one of the sources of calcitriol production. During pregnancy, closer to 12 weeks, the concentration of 1,25(OH)₂D more than doubles, which confirms the important role of vitamin D in supporting gestation and controlling fetal development processes. It regulates decidualization, implantation, and expression of placental lactogen. In addition, vitamin D affects the secretion of progesterone and estradiol, human chorionic gonadotropin (hCG), calcium absorption by the placenta and placental immunomodulation. Thus, active metabolites of vitamin D have been shown to affect numerous physiological processes. Currently, the term vitamin D combines more than 6 molecules of a similar nature, of which 2 molecules of steroid prohormones - D₂ and D₃ - have the greatest biological significance. Vitamins D₂ and D₃ are biologically inert. To convert them into the active form of D-hormone (1,25((OH))₂D), which binds to its receptors in tissues, two successive hydroxylations are necessary. The first occurs in the liver and is converted by the mitochondrial enzyme vitamin D 25-hydroxylase (CYP27A1) to 25-hydroxyvitamin D (25((OH))D), also known as calcidiol. The second hydroxylation occurs in the kidneys, resulting in the synthesis of biologically active 1,25-dihydroxyvitamin D (1,25(OH)₂D), or calcitriol, under the action of mitochondrial 1 α -hydroxylase (CYP27B1). The enzyme 1 α -hydroxylase is also found in many other tissues, which contributes to the local conversion of 25(OH)D to active Vitamin D levels influence 172 key physiological indicators of human health associated with the risk of developing various diseases and the vitamin D receptor (VDR) is found in more than 38 organs and tissues (20,27). There are "classical" effects of D-hormone associated with its influence on calcium-phosphorus metabolism and bone mineral density, and "non-classical" biological effects. The "non-classical" (extraosseous) effects of D-hormone include inhibition of cell proliferation and angiogenesis, stimulation of the production of insulin and cathelicidins (antimicrobial peptides), inhibition of renin production, anti-inflammatory, immunomodulatory, antibacterial, antitumor and a number of other properties.

Similar results were obtained in a study by some co-workers, which indicates that vitamin D in human ovarian granulosa cells in vitro stimulates the production of estradiol, estrone, progesterone, and insulin-like growth factor 1 (19,26). Reports on the prevalence of vitamin D deficiency and insufficiency in various countries around the world show the widespread prevalence of this problem among different population groups, regardless of the standard of living in the country and its geographical location. In the epidermis, cholecalciferol binds to vitamin D-binding protein and 70% of it comes from the bloodstream to the liver, and the other part goes to fat cells, where a vitamin D depot is formed. It has been shown that when human skin is exposed to sunlight at a single erythema dose,

the level of vitamin D3 in the blood increases in the same way as after ingestion of 10,000 IU of vitamin D3 (3,25). However, the development of hypervitaminosis D during prolonged sun exposure does not occur due to blocking the entry of excess vitamin D from the skin into the bloodstream and its transformation into inactive compounds. With age, the content of 7-dehydrocholesterol in the epidermis decreases; accordingly, the synthesis of vitamin D3 decreases and after 65 years its level decreases by more than 4 times. 25-hydroxycholecalciferol is considered the most accurate indicator of vitamin D levels. This is because 25(OH)D has a long half-life (about 3 weeks)(9).

The currently used 500 IU per day is sufficient to maintain the optimal level of metabolism of Ca and P, but is not sufficient to implement the non-calcemic functions of cholecalciferol. The main manifestations of the activity of its metabolites in the body are: regulation of mineral metabolism, especially calcium and phosphorus; regulation of bone growth, remodeling and repair; inhibition of the renin-angiotensin system; anti-inflammatory, regulation of cell growth and differentiation, angiogenesis; regulation of neuromuscular conduction (13,24). When prescribing vitamin D preparations, laboratory monitoring is recommended 2–3 months after the start of treatment. To assess vitamin D status in pregnant women, the total concentration of 25(OH)D [25(OH)D2 and D3] in blood serum was determined using an enzyme immunoassay method (7,23). The results of 25(OH)D levels were assessed in accordance with European recommendations. At a vitamin D level <20 ng/ml, its deficiency was diagnosed; at a concentration of 20–30 ng/ml – deficiency (suboptimal status); 30–100 ng/ml was accepted as the optimal level. To prevent vitamin D deficiency, pregnant and lactating women need to receive at least 800-1,200 IU daily. 90 pregnant women were observed, women received 5,000 units of vitamin D daily until 26 weeks of pregnancy, and the control group (44 patients) received placebo. As a result, women taking placebo had a statistically higher rate of development than those in the study group (35.9% vs. 10.9 P <0.005), indicating the effectiveness of vitamin D supplementation in preventing preterm birth in the first and second trimesters of pregnancy (8,12,22). Summarizing the available data, it can be assumed that women at risk of preeclampsia, gestational diabetes mellitus, recurrent miscarriage, placental insufficiency and IUI should take at least 4000 IU of vitamin D at the preconception stage and from early pregnancy (14). For healthy women with a vitamin D level of 25-30 ng/ml, 2000 IU may be used; above 30 ng/ml, the currently recommended dose is 1000 IU (17,21). Preventing prematurity is a global priority; thus, in the future, women at risk of vitamin D deficiency may be assessed and vitamin D supplements prescribed.

Literature:

1. Alpert P.T., Shaikh U. The effects of vitamin D deficiency and insufficiency on the endocrine and paracrine systems. *Biol Res Nur.* 2007;9(2):117–29.
2. Farhan C, Eleftheria L, Katerina V, Guillermina G Immunomodulatory effects of vitamin D in pregnancy and beyond frontiers In *Immunology* November (2019) 10, 1-17
3. Heaney R.P. Vitamin D in health and disease. *Clin J Am Soc Nephrol.* 2008;3(5):1535–41.
4. Heydena E.L., Wimalawansa S.J. Vitamin D: effects on human reproduction, pregnancy, and fetal Well-Being *Journal Of Steroid Biochemistry And Molecular Biology* 180 (2018) 41–50
5. Holick M.F. Medical progress: vitamin D deficiency. *New Engl J Med.* 2007;357(3):266–81.
6. Ikhtiyarova, G. A. (2022). OUTCOMES OF ASSISTED REPRODUCTIVE TECHNOLOGY PROGRAMS IN PATIENTS WITH A HISTORY OF RELATED PREGNANCY MISSION. *Thematics Journal of Microbiology*, 6(1).
7. Ikhtiyarova, G. A. Diagnostics, treatment and obstetric tactics for fetal loss syndrome. *F: News of Dermatology and reproductive health*, (1.2016), S74-75.

8. Ikhtiyarova, G. A., & Aslanova, M. (2019). Zh., Dustova NK Microbiological changes in pregnant women with antenatal fetal death. *European journal of research*, (2), 102-109.
9. Ikhtiyarova, G. A., & Sh, K. D. Optimization of prenatal screening for diagnostics of intrauterine anomalies of fetal development Journal of ethics and diversity in international communication 2022 Issue 8.
10. Ikhtiyarova, G. A., Aslanova, M. J., & Dustova, N. K. (2020). EARLY MARKERS OF DIAGNOSIS FETAL GROWTH RESTRICTION SYNDROME. *Reproductive Medicine*, (3 (44)), 48-51.
11. Ikhtiyarova, G. A., Dustova, N. K., & Qayumova, G. (2017). Diagnostic characteristics of pregnancy in women with antenatal fetal death. *European Journal of Research*, (5), 5.
12. Ikhtiyarova, G. A., Dustova, N. K., & Tosheva, I. I. (2020). KurbanovaZ. Sh, Navruzova NO "Clinical manifestations of COVID-19 coronavirus infection in pregnant women, measures for pregnancy and childbirth" Methodical recommendation.
13. IKHTIYAROVA, G. A., DUSTOVA, N. K., KHASANOVA, M. A., SULEYMANOVA, G. S., & DAVLATOV, S. S. (2020). Pathomorphological changes of the placenta in pregnant women infected with Coronavirus COVID-19. *International Journal of Pharmaceutical Research* (09752366), 12(3).
14. Karimova, G. K., Navruzova, N. O., & Nurilloeva Sh, N. (2020). An individual approach to the management of gestational diabetes. *European Journal of Molecular & Clinical Medicine*, 7(2), 6284-6291.
15. Lanham-New S.A., Buttriss J.L., Miles L.M. et al. Proceedings of the rank forum on vitamin D. *Br J Nutr*. 2011;105(1):144-56.
16. Lisa M. Bodnar, Mark A. Klebanoff, Alison D. Gemand, Robert W. Platt, W. Tony Parks, Janet M. Catov, And Hyagriv N. Simhan Maternal vitamin D status and spontaneous preterm birth by placental Histology In The Us Collaborative Perinatal Project *Am J Epidemiol*. 2014;179(2):168-176
17. Muxiddinovna, I. M. (2022). Effects of Energy Drinks on Biochemical and Sperm Parameters in Albino Rats. *CENTRAL ASIAN JOURNAL OF MEDICAL AND NATURAL SCIENCES*, 3(3), 126-131
18. Muxiddinovna, I. M. (2022). Effects of Energy Drinks on Biochemical and Sperm Parameters in Albino Rats. *CENTRAL ASIAN JOURNAL OF MEDICAL AND NATURAL SCIENCES*, 3(3), 126-131.
19. Muxiddinovna, I. M. (2022). IMPACT OF ENERGY DRINKS AND THEIR COMBINATION WITH ALCOHOL TO THE RATS METOBOLISM. *Gospodarka i Innovacje*, 22, 544-549.
20. Muxiddinovna, I. M., & Sobirovna, A. Z. (2022). Anemia Iron Deficiency in Pregnancy. *Central Asian Journal of Literature, Philosophy and Culture*, 3(11), 191-199.
21. Muxiddinovna, I. M., & Sobirovna, A. Z. (2022). Pregnancy with Twins with Preeclampsia. *Central Asian Journal of Literature, Philosophy and Culture*, 3(11), 212-221.
22. Narzullaeva, N. S. (2021). Gynecological and somatic history of women with uterine myoma according to retrospective analysis. *Акуш., гинекол., перинатол.*, (2), 86.
23. Razokov, I. B., Rahimov, D. A., Ismatova, M. M., & Kurbonzoda, S. N. MAIN FACTORS CAUSING IRON-DEFICIENT ANEMIA IN CHILDREN OF EARLY AGES. In *SPECIAL EDITION FOR 1st BUKHARA INTERNATIONAL MEDICAL STUDENTS CONFERENCE* (p. 61).

24. Salimova, T. B. (2022). Features of the Course of Pregnancy in Pregnant Women with Fetal Growth Restriction Syndrom and the Role of Doppler Velocimetry. *Central Asian Journal of Medical and Natural Science*, 3(6), 557-563.
25. Абашова Е.И., Ярмолинская М.И., Мишарина Е.В. Роль витамина D в репродукции // Доктор.Ру. 2018. № 10 (154). С. 37–42.
26. Баклейчева М.О., Ковалева И.В., Беспалова О.Н., Коган И.Ю. Влияние витамина D на репродуктивное здоровье женщины // Журнал Акушерства и женских болезней. — 2018. — Т. 67. — № 3. — С. 4–19. Doi: 10.17816/Jowd6734-19 Поступила в редакцию: 21.03.2018
27. Вислоцкий.Н.А,Хабаров.С.В.Международный опыт применения препаратов витамина D с целью профилактики, осложнений беременности и неблагоприятных перинатальных исходов.Вестник Новых медицинских технологий .Вестник Вонгтму.2020.Ст47-52.
28. Герман А. И. Факторы риска преждевременных родов «Актуальные проблемы современной медицины и фармации – 2020» Ст 34-37
29. Горелова И.В., Попова П.В., М.В. Рулёв М.В. Витамин D и репродуктивное здоровье Проблемы Эндокринологии 2020;66(5)Ст:96-101
30. Денисова Т.Г., Грузинова Е.Н., Васильева Э.Н., Сидоров А.Е, Денисова Е.А. Обеспеченность витамином D пациенток с неразвивающейся беременностью Acta Medica Eurasica. 2021. № 1 Ст 11-17
31. Жукова Э., Сычев Д., Чилова Р. И Др. Преждевременные роды: диагностика и лечение // Врач. – 2018; 29 (2): 78–80. Doi: 10.29296/25877305-2018-02-21
32. Ильясов, А. С., & Исматова, М. М. (2022). ЖИНСКИЙ АБЗОЛАРГА ЭНЕРГЕТИК ИЧИМЛИКЛАРНИНГ САЛБИЙ ТАСИРИ. *Uzbek Scholar Journal*, 5, 66-69.
33. Исмоилова, М. З. (2023). Развитие Прогностических Маркер У Женщин Проявляющий Преждевременных Родов С Инфекциями Мочеполовой Системы. *AMALIY VA TIBBIYOT FANLARI ILMIY JURNALI*, 2(5), 362-367.
34. Исмоилова, М. З., & Туксанова, Д. И. (2021). MICROBIOLOGICAL RECOGNITION OF ANTIBODIES TO ANTIGENS OF MICROORGANISMS IN WOMEN WITH INFLAMMATORY DISEASES OF THE GENITAL. *УЗБЕКСКИЙ МЕДИЦИНСКИЙ ЖУРНАЛ*, 2(5)
35. Ихтиярова, Г. А., & Розикова, Д. К. (2023). МИКРОБИОЛОГИЧЕСКИЕ ИЗМЕНЕНИЯ У БЕРЕМЕННЫХ С РЕПРОДУКТИВНЫМИ ПОТЕРЯМИ В АНАМНЕЗЕ. *Finland International Scientific Journal of Education, Social Science & Humanities*, 11(4), 1002-1008.
36. Каримова, Г. К., Ихтиярова, Г. А., & Наврузова, Н. О. (2020). Скрининг диагностика гестационного диабета. *Новый день в медицине*, (1), 220-222.
37. Кузибаева Р. К. Причины и результаты преждевременных родов// В. Н. М. Т. 2015. Т. 22.2. Ст
38. Нарзуллоева, Н. С. (2023). НЕОНАТАЛЬНЫЕ ОСЛОЖНЕНИЯ НА ФОНЕ САХАРНОГО ДИАБЕТА У БЕРЕМЕННЫХ. *European Journal of Interdisciplinary Research and Development*, 15, 333-342.
39. Петров Ю. А., Багновская А. Г., Блесманович А. Е. Влияние микронутриентов на репродуктивное здоровье женщины Журнал гинекология №1 (71) • 2020 Ст.40-43
40. Пигарова Е.А., Рожинская Л.Я., Белая Ж.Е., Дзеранова Л.К, Каронова Т.Л., Ильин А.В., Мельниченко Г.А.,Дедов И.И. Клинические Рекомендации Российской Ассоциации Эндокринологов по диагностике,

лечению и профилактике дефицита витамина D у взрослых Проблемы Эндокринологии, 4, 2016 Ст.60-84.

41. Рамазанова Ф.У., Хамошина М.Б., Гигани О.О., Тулупова М.С. Витамин D и его рецепторы в патогенезе акушерских осложнений: Современный взгляд на проблему // Акушерство И Гинекология: Новости, Мнения, Обучение. 2020. Т. 8, № 3. С. 75–80. Doi: 10.24411/2303-9698-2020-13011
42. Салимова, Т. Б., & Дустова, Н. (2023). ПРИЧИНЫ РИСКА РАЗВИТИЯ СИНДРОМА ОТСТАВАНИЯ РОСТА ПЛОДА В БУХАРСКОЙ ПОПУЛЯЦИИ. *ЖУРНАЛ РЕПРОДУКТИВНОГО ЗДОРОВЬЯ И УРО-НЕФРОЛОГИЧЕСКИХ ИССЛЕДОВАНИЙ*, 4(4).

